

region of the hTR, wherein the first nucleotide sequence within the accessible region is selected from the group consisting of nucleotides 137-196, nucleotides 290-319, and nucleotides 350-380 of hTR, and wherein the polynucleotide comprises a nucleotide analog or a non-naturally occurring nucleotide linkage selected from the group consisting of phosphorothioates, phosphoramidates, methyl phosphonates, chiral-methyl phosphonates, 2-O-methyl ribonucleotides and peptide-nucleic acids.

28. (new) A polynucleotide according to claim 27, when the polynucleotide sequence is selected from the group consisting of:

CGT TCC TCT TCC TGC GGC CTG AAA CGG TGA (SEQ ID NO:2)

CGT TCC TCT TCC TGC GGC CT (SEQ ID NO:3)

CGT TCC TCT TCC (SEQ ID NO:4)

CTG ACA GAG CCC AAC TCT TCG CGG TGG CAG (SEQ ID NO. 5)

CTG ACA GAG CCC AAC TCT TC (SEQ ID NO:6)

CCA ACT CTT CGC GGT GGC AG (SEQ ID NO:7)

GCT CTA GAA TGA ACG GTG GAA GGC GGC AGG (SEQ ID NO:8)

GCT CTA GAA TGA ACG GTG G (SEQ ID NO:9)

GCT CTA GAA TGA ACG (SEQ ID NO:10)

GCT CTA GAA TG (SEQ ID NO:11)

GCT CTA G (SEQ ID NO:12)

CAT TTT TTG TTT GCT CTA GA (SEQ ID NO:13) and

CGG GCC AGC AGC TGA CA (SEQ ID NO:14).

29. (new) A polynucleotide consisting essentially of a sequence selected from the group consisting of:

CGT TCC TCT TCC TGC GGC CTG AAA CGG TGA (SEQ ID NO:2)

CGT TCC TCT TCC TGC GGC CT (SEQ ID NO:3)

CGT TCC TCT TCC (SEQ ID NO:4)

CTG ACA GAG CCC AAC TCT TCG CGG TGG CAG (SEQ ID NO. 5)

CTG ACA GAG CCC AAC TCT TC (SEQ ID NO:6)

CCA ACT CTT CGC GGT GGC AG (SEQ ID NO:7)